

# Carmel Area WD (CA) - Facility Influent


Sample collection date: July 12, 2020

# SARS-CoV-2 virus in sewage

# **DETECTED**

Virus concentration per liter of sewage 17,822 copies

# Relative concentration over time SARS-CoV-2 normalized concentration in blue New cases for your county in black Concentration levels of SARS-CoV-2 at your facility over time, compared with new case trends for your county. Data is normalized to a fecal marker Change since 175% May Jun Jul Aug Sep Oct






# **About the data**

Our methods for detecting SARS-CoV-2 in sewage are adapted from CDC protocols and available at www.biobot.io/covid19. Our approach relies on detecting genetic fragments of the virus that are excreted in stool, which does not determine if the virus is dead or active.

# Relative concentrations trend with new cases

Concentration levels are trending with the county's new reported cases at the time of sample collection.


This sample in red, all other samples since May in blue.



# Relative concentrations in comparison

Your sample in red, median in black, all other samples collected since May in blue

Your sample has higher concentration levels than **7%** of all quantifiable samples collected since May.



Contextualize SARS-CoV-2 levels in your facility's catchment with other participating facilities.

# **Case Estimates Reflect Active R&D**

Evaluate as beta results when comparing to reported Covid19 cases

For more information, https://doi.org/10.1101/2020.04.05.20051540 read the whitepapers: https://doi.org/10.1101/2020.06.15.20117747

# Reported COVID19 cases in Monterey County, CA

New cases

Cumulative cases

2,419

On July 12, 2020, as reported by USAfacts.org

# Biobot COVID19 case estimate

# 110 cases

(0.77% incidence rate)

Using a reported flow rate of 1.2 MGD

COVID19 sewage sampling Report provided: July 17, 2020

# Biobot's wastewater testing protocols

We are continuously working to improve our protocols to increase the sensitivity of our measurements and reduce variability. We use a qPCR-based method to detect and quantify the SARS-CoV-2 virus and an associated fecal normalization control.

The RNA extraction method changed between protocols v1 and v2 to improve the sensitivity of our measurements. Protocols v2.1-v2.3 reflect changes in our quantification method to further improve the precision of our analysis and reduce variability. Internal tests have been conducted to assure data integrity. The SARS-CoV-2 viral titers that we report may be affect by slight batch effects between these protocol versions.

The sensitivity of our assay varies slightly between these protocols. The exact limit of detection (LOD) that applies to your sample can vary. Generally, the LOD for each protocol is: v1: 6,400 copies/L; v2: 34,000 copies/L; v2.1: 1,700 copies/L; v2.2: 2,100 copies/L; v2.3: 3,600 copies/L.

## Why normalize with a fecal marker?

We normalize SARS-CoV-2 viral titers to account for differences in the total people contributing to each sample. We use PMMV as this internal control, which is an RNA virus that is excreted in stool.

### Biobot's COVID19 case estimate

We measure the concentration of SARS-CoV-2 in sewage. We convert our measurements into a COVID19 case estimate using the following basic equation:

# Number of infected people =

(total amount of virus per day) / (virus shed per infected person per day)

## Comparing with confirmed clinical cases

Our COVID19 case estimates may not match the confirmed case numbers in the community for a variety of reasons. Clinical testing may not represent the entire infected population. The SARS-CoV-2 virus may start being shed before patients develop symptoms and seek testing.

We encourage you to share these reports with your local public health officials to compare our case estimates with the number of confirmed cases in your local community.

### Data use

The Biobot COVID19 case estimates provide an alternative metric to guide responses to the outbreak. We recommend that you share this information with local public health officials. We believe this work will have the greatest impact on a statewide level, and hope that you will reach out to your state officials and encourage the expansion of our partnership across your state.

# Pioneering a better way to track Covid-19

We are the nation's leader in wastewater epidemiology. We analyze sewage to determine the presence of infected individuals and estimate the number of Covid-19 cases, providing:

- <u>Trend analysis</u>: Determine when to safely re-open, and keep open, our communities and businesses through frequent sampling.
- <u>Early warning</u>: Detect the re-emergence of Covid-19 in a population to rapidly take action and contain new outbreaks.
- <u>Testing at scale</u>: For a fraction of the cost of mass individual testing, get an overview of the scope of the outbreak.

For questions, email support@biobot.io